



Borregaard
LignoTech

LignoTech USA, Inc.
100 Grand Avenue
Rothschild, WI 54474
Tel: (715)-355-6544
Fax: (715)-355-3674

BorreGRO® HA-2 **(Powder)**

Humic Acid

GUIDELINES FOR USE:

BorreGRO® HA-2 is potassium humate.

BorreGRO® HA-2 promotes the uptake of micronutrients.

BorreGRO® HA-2 may be applied alone or with liquid fertilizers with a pH 7.0 or higher.

Compatibility can be simply checked by:

- Mixing each product in a clear container in the same proportion to be used in the field
- Agitating vigorously for at least 30 seconds
- Visually observing for precipitate

To prepare a 6% aqueous solution of **BorreGRO® HA-2**, dissolve one pound of powder in one gallon of water.

“NOT A PLANT FOOD INGREDIENT”

GUARANTEED ANALYSIS

60% Humic Acid derived from Leonardite

- WARNING: NUISANCE DUST
- AVOID HIGH CONCENTRATIONS
 - MINIMIZE CONTACT WITH EYES, SKIN & CLOTHING
 - KEEP AWAY FROM HEAT, SPARKS & OPEN FLAME

LignoTech USA, Inc. does not warrant or guarantee in any manner whatsoever, including the warranty of merchantability or fitness for the end user the accuracy of the information and procedures listed herein and will not be responsible for any damage resulting from their use.

Contact your LignoTech Sales Representative for additional information.

CA 2956

Net Weight: _____

Lot Number: _____

BorreGRO® HA-2 can be applied to all crops including fruits, vegetables, trees, vines, flowers, ornamentals, turf, grain and other row crops.

Application rates and frequency are dependent upon soil and climate conditions present. Typical application rates range from 1/2 to 10 pounds per crop per acre per season.

BorreGRO® HA-2 may be applied through conventional methods but not limited to foliar spray, soil application, water run, shanking in or hydroponically.

Follow the labeled rates for nutrients and/or consult with your local PCA.