



VISTA CLEAR SPRAY OIL II

 For Organic Production

For Use in Selected Berries, Cotton, Field and Sweet Corn, Selected Fruit Trees, Grapes, Grasses Grown for Seed, Hops, Spearmint, Selected Nut Trees, Sugar Beets, Tobacco, Ornamentals, and Selected Vegetable Crops

Controls Fungal Diseases, Insects, Mites, and Virus-Transmitting Aphids

Active Ingredient:

Mineral Oil ¹	98.4%
Other Ingredients	1.6%
TOTAL	100.0%

¹Contains petroleum distillates

Minimum Unsulfonated Residue.....98%

KEEP OUT OF REACH OF CHILDREN

CAUTION

See inside for First Aid, Precautionary Statements, Directions for Use, Conditions of Sale and Warranty, and state-specific crop and/or use site restrictions.

EPA Reg. No. 90784-5
EPA Est. No. 041696-CA-004
EPA Est. No. 090784-CA-001
EPA Est. No. 066663-CA-001



Mar Vista
Resources®
Providing Resources for Value Creation™



Guaranteed, Manufactured & Formulated By: Mar Vista Resources, LLC at 745 North Ave • Corcoran, CA 93212 • Office: (559) 992-4535

FIRST AID

- | | |
|-------------------------------|---|
| If Swallowed | <ul style="list-style-type: none">• Immediately call a poison control center or doctor.• Do not induce vomiting unless told to do so by a poison control center or doctor.• Do not give any liquid to the person.• Do not give anything by mouth to an unconscious person. |
| If Inhaled | <ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.• Call a poison control center or doctor for further treatment advice. |
| If in Eyes | <ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice. |
| If on Skin or Clothing | <ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice. |

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also call National Poison Control Center at 1-800-222-1222 for emergency medical treatment information.

NOTE TO PHYSICIAN

Contains petroleum distillate. Vomiting may cause aspiration pneumonia.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Harmful if swallowed. If swallowed do not induce vomiting. Call a physician. Avoid breathing vapors and spray mists.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of barrier laminate, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, or viton ≥ 14 mils
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with the product's concentrate. Do not reuse them.

Engineering Controls:

Pilots must use an enclosed cockpit that meets the requirements listed in the WPS for agricultural pesticides (40 CFR 170.240(d)(6)).

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Then wash the outside of gloves before removing them. Wash thoroughly and change into clean clothing as soon as possible.

Environmental Hazards

This product is toxic to fish. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements, specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow entry into treated areas during the restricted entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves, made of barrier laminate, nitrile rubber \geq 14 mils, neoprene rubber \geq 14 mils, or viton \geq 14 mils
- Shoes plus socks

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Keep away from heat, sparks and open flames. Keep container closed when not in use to prevent entry of water.

Pesticide Disposal: Open dumping is prohibited. Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Handling:

Nonrefillable Container (rigid material; ≤ 5 gallons): Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling, if available, or dispose of empty container in a sanitary landfill or by other procedures allowed by state and local authorities.

Nonrefillable Containers (rigid material; > 5 gallons): Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Offer for recycling, if available, or dispose of empty container in a sanitary landfill or by other procedures allowed by state and local authorities.

Refillable Container > 5 gallons: Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Offer for recycling, if available, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

FOR 24-HOUR EMERGENCY ASSISTANCE (SPILL, LEAK, OR FIRE), CALL INFOTRAC AT 1-800-535-5053.

PRODUCT INFORMATION

This product controls select fungal diseases, mites, and insects, including aphids which transmit plant viruses. The product works by smothering the target pest, so thorough coverage is required for best control. This product does not provide residual pest control.

Some crop varieties are sensitive to oil sprays. Growers inexperienced with applying this product at a rate, spray interval, or in tank mix should apply to a small area of the crop using the recommended nozzles, spray pressure, and tractor speed. Wait several days to determine if phytotoxicity occurs before applying the mixture to large acreages.

APPLICATION RATE

Unless directed otherwise in the crop-specific sections of this label, use the application rates in this section. When a range of application rates is allowed, use the higher rate when mite or insect infestation is heavy or disease pressure is high.

Thoroughly cover the plants, using from 25 to 150 gallons of spray solution per acre (depending on plant size) and a tractor speed of 3½ to 4 mph. Vary the amount of spray solution applied by increasing or decreasing the number of nozzles used rather than changing nozzle size, spray pressure, or sprayer speed.

APPLICATION TIMING

Unless directed otherwise in the crop-specific sections of this label, use the application timing in this section. When varying application timing is allowed, use the shorter spray interval when insect or mite infestation is heavy or disease pressure is high.

Insecticide/Miticide Use: Begin spraying as soon as the plants emerge. Thereafter, spray once or twice weekly, depending on the level of infestation. Continue application through harvest, as needed. See the exception for table grapes in the crop-specific portion of this label.

Insecticide/Fungicide Use: Spray weekly or once every two weeks, depending on disease severity.

Aphid Control to Reduce Plant Virus Transmission: Begin sprays when winged aphids appear. Thereafter, spray once or twice weekly, depending on the level of aphid or virus pressure. See the specific instructions for cucurbits in the Vegetable portion of this label.

EQUIPMENT

For best control of insects, mites, and specified diseases, thorough coverage is essential. To ensure the best coverage, apply by ground using the specified dilution, proper spray pressure, nozzles (generally hollow cone), nozzle spacing, and sprayer speed.

Plant Virus Use: Use ceramic hollow cone nozzles and spray at no less than 400 psi.

Insecticide/Fungicide Use: Spray to cover the upper leaf surface. Although not required, using an application system for vegetables/plant virus results in maximum coverage and least potential for phytotoxicity. Do not apply via chemigation.

SPRAY DRIFT MANAGEMENT

A variety of factors including weather conditions (e.g., wind direction, wind speed, temperature, and relative humidity) and method of application (ground, aerial, or airblast) can influence pesticide drift. The applicator and grower must evaluate all factors and make appropriate adjustments when applying this product.

Wind Speed

Do not apply at wind speeds greater than 15 mph at the application site.

Temperature Inversions

If applying at wind speeds less than 3 mph, the applicator must determine if a) conditions of temperature inversion exist, or b) stable atmospheric conditions exist at or below nozzle height. Do not make applications into areas of temperature inversions or stable atmospheric conditions.

Droplet Size

Apply as a medium or coarser spray (ASABE standard 572).

Ground Boom Applications

For ground boom applications, apply with nozzle height no more than 4 feet above the ground or crop canopy and when wind speed is 10 mph or less at the application site as measured by an anemometer. Apply as a medium or

coarser spray according to ASABE standard 572 definition for standard nozzles or minimum volume mean diameter (VMD) for spinning atomizer nozzles.

Airblast Applications

For orchard/vineyard airblast applications, do not direct spray above trees/vines and turn off outward pointing nozzles at row ends and outer rows. To minimize spray loss over the top in orchard applications, direct spray into the canopy. Apply only when wind speed is 3 -10 mph at the application site as measured by an anemometer outside of the orchard/vineyard on the upwind side.

Aerial Applications

Do not release spray at a height greater than 10 feet above the ground, top of crop, or above the orchard canopy.

Boom Length

The boom length must not exceed 75% of the wingspan or 90% of the rotor blade's diameter.

Swath Adjustment

When applications are made with a crosswind, the swath will be displaced downwind. The applicator must compensate for this displacement at the downward edge of the application area by adjusting the path of the aircraft upwind. Leave at least one swath unsprayed at the downwind edge of the treated field.

Wind Speed

Do not apply at wind speeds greater than 15 mph at the application site.

Droplet Size

Apply as a medium or coarser spray (ASABE standard 572), and the minimum volume mean diameter (VMD) for spinning atomizer nozzles.

RESTRICTIONS and LIMITATIONS for ALL CROPS

Also, see restrictions and limitations in crop-specific instructions

This product may be used up to the day of harvest for all crops.

DO NOT apply through any type of irrigation system.

DO NOT apply during or immediately prior to temperatures over 95° F, or when the temperature approaches freezing, or during hot, dry winds, rain or other unsuitable conditions.

DO NOT overspray or double spray.

Plant should be in vigorous condition, not under stress from soil moisture, disease, insects or any other conditions unfavorable to plant growth.

CHEMICAL AND PHYSICAL COMPATIBILITY AND PHYTOTOXICITY

Add the water carrier and all other ingredients to the spray tank before adding this product. Perform a mixing test to check the compatibility of this product with the tank mix partner(s) before adding any other product to the spray tank. This product has not been tested on all species or varieties. Before treating a large area, treat a small area and observe prior to full scale application.

TANK-MIXING

This product may be tank mixed with one or more pesticide products registered for use in each crop, per the instructions in this label and the respective product labels. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

DO NOT tank mix this product with spreader stickers or pinolene based products. Wait at least 10 days between an oil application and spraying pinolene-based products with fruit present.

DO NOT tank mix with highly ionized nutrient materials.

DO NOT apply sulfur within 10 days of an oil application on grapes. On all other crops, do not apply sulfur within 10 days of an oil application and do not apply oil within 14 days of an application of wettable or dusting sulfur.

DO NOT use captan or folpet with or following an oil spray.

DO NOT use chlorothalonil, dicloran, or permethrin within 14 days following an oil application.

DO NOT use chlorothalonil or dimethoate in a spray program with oil on grapes, ornamentals and strawberries.

DO NOT use propargite with an oil spray or within 30 days before or after an oil spray.

FIELD USE

Apply as directed in crop-specific sections for both bearing and non-bearing crops, except for apples, cherries, peaches and pears. See separate use directions for bearing apples, cherries, peaches, and pears and non-bearing apples, cherries, peaches, and pears.

CROP-SPECIFIC USE DIRECTIONS

Crops	Target Diseases	Use Rate/100 gal. Water	Application Timing and Restrictions
Almonds Apricots	Scales, Mites Fruit Tree Leafroller*	1-1 ½ gals 2-3gals	Dormant or Delayed Dormant Use
	Scales, Mites	1 gals	Summer (foliar or cover sprays) and Post harvest Do Not apply to oil sensitive varieties.
Apples	Apple Red Bug European Fruit Lecanium Scale San Jose Scale Fruit Tree Leafroller* European Red Mites	1-2 gals	Dormant or Delayed Dormant to ½ Green Tip Use
	Scurfy Scales	3 gals	
	Mites (Including European Red Mite) Rust Mite*, Leafhopper* San Jose Scale Rosy Apple Aphid* Fruit Tree Leafroller*	1-2 gals	Summer Foliar or Cover Spray Use. Apply every 10-14 days depending on the level of pest pressure. Postharvest use to reduce overwintering pest population. Do Not apply to oil sensitive varieties.
	Powdery Mildew	1-2 gals	Apply at tight cluster and continue every 10-14 days through second cover spray. Use the higher rate and shorter treatment interval when disease pressure is high.
Blueberries Bushberries Caneberries	Mites Powdery Mildew Rust	¾ to 1½ gals	Spray using ground equipment and ensure thorough coverage of the foliage by using a spray pressure of at least 200 psi.

* Asterisk denotes use on crop or pest not approved in California

Crops	Target Diseases	Use Rate/100 gal. Water	Application Timing and Restrictions
Cherries (bearing)	Fruit Tree Leafroller*	1-2 gals	Dormant or Delayed Dormant Use
	Mites, Scale	1-2 gals	Summer Foliar Use. Apply when mites first appear. Repeat every 10-14 days. Postharvest sprays may be made to maintain control and reduce overwintering pest population. Do Not apply to oil sensitive varieties.
	Powdery Mildew	1-2 gals	Preharvest Use: Use up to pit hardening stage. Make application every 10 to 14 days. Use the higher rate and shorter treatment interval when disease pressure is high.
Cotton Field and Sweet Corn Sugar Beets Tobacco	Earthworm Fall armyworm Corn rootworm Leafhopper Leafminer Mites Whiteflies Cercospora	¾ to 1½ gals	Using ground equipment, ensure thorough coverage of the foliage. To control Cercospora use a minimum of 200 psi spray pressure.
Grapes	Mealybugs Mite Leafhopper* Whiteflies*	1 to 2 gals	Apply using ground equipment and ensuring thorough coverage of leaf surface every 10-14 days.
	Powdery Mildew Botrytis bunch rot		To control powdery mildew, apply this product pre-bloom and every 10 to 21 days thereafter, depending on disease pressure. Use the higher rate and shorter spray interval when powdery mildew pressure is high. To control Botrytis bunch rot, apply at bloom and repeat before bunch-closing and veraison. A final application may be made at harvest, if needed. Use the higher rate and shorter spray interval when Botrytis bunch rot is severe.
<p>RESTRICTIONS AND LIMITATIONS: DO NOT apply more than 2 gallons of this product per acre per application. DO NOT apply more than 3 gallons of any mineral spray oil per acre per application. Wait AT LEAST 10 days between applications of sulfur products and this product. DO NOT use products containing chlorothalonil or dimethoate during the same growing season as this product. When the crop is fruiting, DO NOT use copper products and this product together. DO NOT apply within 14 days of harvesting table grapes.</p>			

* Asterisk denotes use on crop or pest not approved in California

Crops	Target Diseases	Use Rate/100 gal. Water	Application Timing and Restrictions
Hops*	Mites Powdery Mildew	1 to 2 gals	Apply every 10 to 14 days. For powdery mildew control, begin spraying at early leaf stage.
	RESTRICTION AND LIMITATIONS: DO NOT apply Folpet during the same growing season as this product. DO NOT spray this product after burrs begin to develop.		
Mango*	Mites	1-1½ gals	Apply every 14-21 days.
	Powdery Mildew		Begin application before bloom. Repeat spray at bloom and every 14 to 21 days thereafter as needed.
Nectarines Peaches (bearing)	Fruit Tree Leafroller* Cottony Peach Scales, Mites, Scale	2½ gals 3 gals 1 to 2 gals	Dormant or Delayed Dormant Use
	Mites, Scales	1 to 2 gals	Summer Foliar Use. Postharvest application can be made to maintain control and reduce over-wintering pressure.
	Powdery Mildew	1 to 2 gals	To control powdery mildew, begin spraying at 5% bloom. Continue spraying at 10 to 14-day intervals until terminal growth phase stops. Do Not apply to oil sensitive varieties.
Non-bearing Fruit Trees: Apples, Cherries, Nectarines, Peaches, and Pears	Mites Powdery Mildew	1-2 gals	To control mites and powdery mildew, apply at 10 to 14-day intervals. For powdery mildew, use 200 psi spray pressure and apply before the disease is present; thoroughly cover leaf surface. Use the higher rate and shorter spray intervals under heavy pest pressure.
Olives	Mites, Scales	1½ gals	Apply during prebloom to postbloom buckshot stage.
Papaya*	Mites, Powdery Mildew, Papaya ringspot virus	¾ to 1½ gals	To control mites and powdery mildew spray every 10 to 14 days. Use the higher rate and shorter spray interval when pest pressure is high. To control aphids bearing papaya ringspot virus, begin application when the seeds are germinated. Then spray every week using hollow cone nozzle and 400 psi spray pressure.

* Asterisk denotes use on crop or pest not approved in California

Crops	Target Diseases	Use Rate/100 gal. Water	Application Timing and Restrictions
Pears (bearing)	Fruit tree leafroller* Pear leaf blister mite Pear psylla*	3 gals 2 gals	Dormant or Delayed Dormant Use, up to and including Petal Fall.
	Red spider mite San Jose & Italian pear scale (Summer forms) Leafhopper* Pear psylla*	1 gal 1 to 1½ gals 1 to 1½ gals 1½ to 2 gals	Summer Use (foliar or cover sprays). Postharvest Use. To reduce overwintering populations. Do Not apply to oil sensitive varieties.
	Powdery Mildew	1 to 2 gals	To control powdery mildew, apply at bud burst. Continue every 10 to 14 days, using the higher rate and shorter spray interval when disease pressure is high.
Pecans	Obscure scale	3 gals	Dormant Spray
Plums Prunes	European red mites European fruit lecanium scale Mites, Scale	2 gals 2 gals 1 to 1½ gals	Dormant to ½ green tip
	Mites, Scales	1 to 1½ gals	Summer Foliar Use
Strawberries	Leafminers* Mites Botrytis Powdery Mildew	¾ gals	Use hollow cone spray nozzles and at least 400 psi.
	RESTRICTIONS AND LIMITATIONS: DO NOT use products containing chlorothalonil or dimethoate during the same growing season as this product.		

* Asterisk denotes use on crop or pest not approved in California

Crops	Target Diseases	Use Rate/100 gal. water	Application Timing and Restrictions
Vegetables: Asparagus, Bean, Beet, Cabbage, Cantaloupe, Casaba, Cauliflower, Celery, Chayote fruit, Chinese Wax gourd, Citron melon, Crenshaw melon, Cucumber, Cucuzza, Eggplant, Gherkin, Golden Pershaw melon, Honey balls, Honeydew melon, Hyotan, Lettuce, Mango melon, Pepper, Persian melon, Potato, Pumpkin, Summer Squash, Tomato Watermelon, Winter Squash	Leafhopper* Leafminers Mites Whiteflies Alternaria Leafspot Gummy Stem Blight Powdery Mildew Rust	$\frac{3}{4}$ to 1½ gals	Using ground equipment, ensure thorough coverage of the upper leaf surface. Use hollow cone nozzles and 400 psi pressure for aphid transmitted plant viruses and insect control. Use at least 200 psi spray pressure for fungal diseases. Spray when winged aphids begin to appear. Spray once per week or twice per week if virus or aphid pressure is high.
	Aphids Transmitting: Cucumcumber Mosaic Virus Papaya Ringspot Virus Pepper Mottle Virus Potato Leafroll Virus Potato Virus Y Tobacco Etch Virus Watermelon Mosaic Virus 2 Zucchini Yellow Mosaic Virus	$\frac{3}{4}$ gal	
	In cucurbit crops (cantaloupe, casaba, Chayote fruit, Chinese wax gourd, citron melon, crenshaw melon, cucumber, cucuzza, gherkin, golden pershaw melon, honey balls, honeydew melon, hyotan, mango melon, Persian melon, pumpkin, summer squash, watermelon, and winter squash) begin applications when 50% of the plants have germinated. When cucurbits are growing rapidly, spray twice per week. RESTRICTION AND LIMITATIONS: DO NOT apply this product unless temperatures are 50°F or higher.		
Walnuts	Mites, Scale	1 to 1½ gals	Delayed Dormant Use
	Mites, Scale	$\frac{1}{2}$ to 1 gal	Spring to Mid-Summer Use
	RESTRICTIONS AND LIMITATIONS: DO NOT spray this product after husk split.		

* Asterisk denotes use on crop or pest not approved in California

Crops	Target Diseases	Use Rate	Application Timing
Spearmint	Mites Powdery Mildew	1 to 2 gals/A with at least 20 gal/A of water	For Powdery mildew, apply by ground when disease is detected or when mint is 5 to 6 inches tall ensure thorough coverage of foliage. Repeat treatment if needed, at 14 to 21-day intervals. Use higher rate and shorter treatment interval when disease pressure is high.

Crops	Target Diseases	Use Rate/1 gal water	Application Timing
Ornamentals: Chrysanthemum, Philodendron, Poinsettia	Leafminers, Mealybugs, Mites, Scale Insects, Whiteflies	1 oz.	To control the listed insects, mites, and diseases apply every 7 to 14 days, as needed. Spray to the point of runoff, using hollow cone nozzles and 20 to 100 gallons of the dilution (depending on plant size). Use the higher rate and shorter spray interval when disease pressure is high.
Rose	Black Spot, Powdery Mildew	1 to 2 oz.	
<p>Vary the amount of spray solution applied by increasing or decreasing the number of nozzles used rather than changing nozzle size, spray pressure, or sprayer speed.</p> <p>RESTRICTIONS AND LIMITATIONS: DO NOT use products containing chlorothalonil or dimethoate during the same growing season as this product.</p>			

CONDITIONS OF SALE AND WARRANTY

NOTICE: The statements made on this label are believed to be true and accurate but because of conditions of use which are beyond our control, Mar Vista Resources, LLC. does not make nor does it authorize any agent or representative to make any warranty, guarantee or representation, expressed or implied, concerning this material or the use thereof, except in conformity with the statements of the label. To the extent permitted by applicable law, neither Mar Vista Resources, LLC. nor the seller shall be held responsible in any manner for any personal injury or property damage or loss resulting to the buyer or to the other person from handling, storage or use of this material not in accordance with the directions. To the extent permitted by applicable law, the buyer assumes all risk and liability resulting from improper handling, storage or use and accepts and uses this material on these conditions.



VISTA CLEAR SPRAY OIL II

For Organic Production

For Use in Selected Berries, Cotton, Field and Sweet Corn, Selected Fruit Trees, Grapes, Grasses Grown for Seed, Hops, Spearmint, Selected Nut Trees, Sugar Beets, Tobacco, Ornamentals, and Selected Vegetable Crops

Controls Fungal Diseases, Insects, Mites, and Virus-Transmitting Aphids

Active Ingredient:

Mineral Oil ¹	98.4%
Other Ingredients	1.6%
TOTAL	100.0%

¹Contains petroleum distillates

Minimum Unsulfonated Residue.....98%

KEEP OUT OF REACH OF CHILDREN

CAUTION

See inside for First Aid, Precautionary Statements, Directions for Use, Conditions of Sale and Warranty, and state-specific crop and/or use site restrictions.

EPA Reg. No. 90784-5
 EPA Est. No. 041696-CA-004
 EPA Est. No. 090784-CA-001
 EPA Est. No. 066663-CA-001

Guaranteed, Manufactured & Formulated by:



Mar Vista Resources, LLC at 745 North Ave • Corcoran, CA 93212 • Office: (559) 992-4535

- Net 1 Gal**
(3.78 L)
- Net 2.5 Gal**
(9.5 L)
- Net 15 Gal**
(57 L)
- Net 275 Gal**
(1,040.9 L)
- Bulk** _____

PULL HERE TO OPEN AND PRESS TO RESEAL ▲