

Bio-Sweet[®]

BIO-FERTILIZER

AGRI LINE



0-13-5

GUARANTEED ANALYSIS:

Available Phosphate (P₂O₅)	13.00%
Soluble Potash (K₂O)	5.00%
Magnesium (Mg)	0.50%
0.50% Water Soluble Magnesium (Mg)	
Sulfur (S)	1.00%
0.05% Water Soluble Copper (Cu)	
Boron (B)	0.05%
Copper (Cu)	0.05%
0.05% Water Soluble Copper (Cu)	
Zinc (Zn)	0.15%
0.05% Water Soluble Zinc (Zn)	
*Maximum Chlorine (Cl) 0.54%	

Derived From: Boron Ethanolamine, Copper-IDHA, Magnesium-IDHA, Phosphoric Acid, Potassium Chloride, Monopotassium Phosphate, Zinc-IDHA.

ALSO CONTAINS NON-PLANTFOOD INGREDIENT(S)
100% TOTAL INERT INGREDIENT(S) (inert non-plant food ingredient) Oligosaccharides (microbial food)

Information regarding the contents and levels of metals in this product is available online at <http://www.aapfo.org/metals.html>

Disclaimer or Warranties: To the extent consistent with applicable law, GROPRO Inc. makes no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond the statements made on this label.

No agent of GROPRO Inc. is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, GROPRO Inc. disclaims any liability whatsoever for special, incidental or consequential damages resulting from the use or handling of this product.

Limitations of Liability: To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use or handling of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid or at GROPRO Inc's election, the replacement of product

Contains Boron. Do not use on plants sensitive to Boron. Use of Boron on plants other than those recommended may result in serious injury to the plants.

PRECAUTIONARY STATEMENTS

Do not get in eyes, on skin or on clothing. Wash hands, forearms, and other exposed areas thoroughly after handling. Do not eat, drink or smoke when using this product. Do not breathe vapors, mist, or spray

CAUTION
KEEP OUT OF REACH OF CHILDREN



May be harmful if swallowed.
Causes mild skin and eye irritation.

How can we help?
1-833-476-7761



MANUFACTURED BY

900 128th Street W,
Burnsville MN 55337
Phone: 1-833-476-7761

www.groproag.com

2.5 Gal (9.46353 L)

10.36 lbs. per gallon @ 68 F
FLORIDA FERTILIZER LICENSE NUMBER F003514

1 A, B, C, D, Root and Tuber Vegetables
Dosage per acre per application: 1-8 pints. Apply within 8 weeks of harvest to correct coloring to desired brix.

2 Leaves of Root and Tuber Vegetables
Dosage per acre per application: 1-8 pints. Apply within 8 weeks of harvest to correct coloring to desired brix.

3-07 A, B, Bulb Vegetable Group
Dosage per acre per application: 1-8 pints. Apply within 8 weeks of harvest to correct coloring to desired brix.

6-22 A, B, C, D, E, F Legume Vegetable Group
Dosage per acre per application: 1-8 pints. Apply within 8 weeks of harvest to correct coloring to desired brix.

8-10 A, B, C Fruiting Vegetable Group
Dosage per acre per application: 1-8 pints. Apply within 8 weeks of harvest to correct coloring to desired brix.

9 A, B Cucurbit Vegetables
Dosage per acre per application: 1-8 pints. Apply within 8 weeks of harvest to correct coloring to desired brix.

10-10 A, B, C Citrus Fruit Group
Dosage per acre per application: 1-8 pints. Apply within 8 weeks of harvest to correct coloring to desired brix.

11-10 Pome Fruit Group
Dosage per acre per application: 1-8 pints. Apply within 8 weeks of harvest to correct coloring to desired brix.

12-12 A, B, C Stone Fruit Group
Dosage per acre per application: 1-8 pints. Apply within 8 weeks of harvest to correct coloring to desired brix.

13-07 A, B, C, D, E, F, G, H, Berry and Small Fruit
Dosage per acre per application: 1-8 pints. Apply within 8 weeks of harvest to correct coloring to desired brix.

All Other Crops not listed that need brix support Bio-Sweet can be applied at a rate of 1-8 pints per acre per application. Apply within 8 weeks of harvest to correct to desired brix.

FIRST AID

• Call a POISON CENTER or doctor immediately. • Rinse mouth. DO NOT induce vomiting. • Do not give anything by mouth to an unconscious person. • Immediately call a POISON CENTER or doctor.

IF IN EYES:

• Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. • Call a poison control center or doctor for treatment advice.

IF INHALED:

• Move person to fresh air and keep at rest in a position comfortable for breathing if they feel unwell. • If not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a POISON CENTER or doctor for treatment advice.

IF ON SKIN OR HAIR:

• Take off immediately all contaminated clothing. Rinse skin with water or shower. • Get medical attention if irritation develops or persists. • Wash contaminated clothing before reuse.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also call 1-800-424-9300 for emergency medical treatment information.